

Baryon magnetic moments in the external field method *

F.X. Lee^a, R. Kelly^a, L. Zhou^a, W. Wilcox^b

^aCenter for Nuclear Studies, Department of Physics, The George Washington University, Washington, DC 20052, USA

^bDepartment of Physics, Baylor University, Waco, TX 76798, USA

We present a calculation of the magnetic moments of the baryon octet and decuplet using the external field method and standard Wilson gauge and fermion actions in the quenched approximation. Progressively smaller static magnetic fields are introduced on a 24^4 lattice at $\beta=6.0$ and the pion mass is probed down to about 500 MeV. Magnetic moments are extracted from the linear response of the masses to the external field.

1. Method

Magnetic moments are an important fundamental property of particles. They determine dynamical response of a bound system to external perturbations, and provide valuable insight into internal strong interaction structure. There are at least two ways to extract the magnetic moment. One is from form factors (three-point functions) where an extrapolation to $G_M(Q^2 = 0)$ is required. The other is the external field method using only two-point functions [1,2,3,4]. Here we report a calculation in the latter. It is an extension of our work on electric and magnetic polarizabilities [5,6,7].

In order to place a magnetic field on the lattice, we construct an analogy to the continuum case. The fermion action is modified by the minimal coupling prescription $D_\mu = \partial_\mu + gG_\mu + qA_\mu$ where q is the charge of the fermion field and A_μ is the vector potential describing the external field. On the lattice, the prescription amounts to a modified link variable $U'_\mu = U_\mu U_\mu^{(B)}$. Choosing $A_y = Bx$, a constant magnetic field B can be introduced in the z -direction. Then the phase factor is $U_\mu^{(B)} = \exp(iqa^2 Bx)$. On a finite lattice with periodic boundary conditions, to get a constant magnetic field, B has to be quantized by the condition $qBa^2 = \frac{2\pi n}{N_x N_y}$, $n = 1, 2, 3, \dots$ to ensure that the magnetic flux through plaquettes in the xy -plane is constant. However, for $N_x = N_y = 16$

and $1/a = 2\text{GeV}$, the lowest field would give the proton a mass shift of about 390 MeV, which is too large (the proton is severely distorted). We abandon the quantization condition and choose to work with smaller fields. To minimize the boundary effects, we work with fixed (or Dirichlet) b.c. in the x -direction and large N_x , so that quarks originating in the middle of the lattice have little chance of propagating to the edge.

We use 24^4 lattice at $\beta = 6.0$, and six kappa values $\kappa=0.1515, 0.1535, 0.1545, 0.1555$, corresponding to pion mass of 1015, 908, 794, 732, 667, 522 MeV. The strange quark mass is set at $\kappa=0.1535$. The source location is $(x,y,z,t)=(12,1,1,2)$. We analyzed 87 configurations. The following five dimensionless numbers $\eta = qBa^2 = +0.00036, -0.00072, +0.00144, -0.00288, +0.00576$ give four small B fields at $eBa^2 = -0.00108, +0.00216, -0.00432, +0.00864$ for both u and d (or s) quarks. Note that "small" here is in the sense that the mass shift is only a fraction of the proton mass: $\mu B/m \sim 0.6$ to 4.8% at the smallest pion mass. In absolute terms, $B \sim 10^{13}$ tesla.

The mass shift in the presence of small fields can be expanded as a polynomial in B ,

$$\delta m(B) = m(B) - m(0) = c_1 B + c_2 B^2 + c_3 B^3 + \dots (1)$$

To eliminate the contamination from the even-power terms, we calculate mass shifts both in the field B and its reverse $-B$, then take the difference and divide by 2. So our calculation is

*presented by F.X. Lee at Lattice 2004.

equivalent to 11 separate spectrum calculations: 5 original η values, 5 reversed, plus the one at zero field to set the baseline. For a Dirac particle of spin s in uniform fields, $E_{\pm} = m_{\pm} \pm \mu B$ where the upper sign means spin up and the lower sign means spin-down, and $\mu = g \frac{e}{2m} s$. We tried the following three methods to extract the g factors, and found they are equivalent within statistical errors.

$$g = \left(\frac{2m_+m_-}{m_+ + m_-} \right) \frac{(E_+ - m_+) - (E_- - m_-)}{eBs} \quad (2)$$

$$g = \frac{m_+(E_+ - m_+) - m_-(E_- - m_-)}{eBs} \quad (3)$$

$$g = \frac{(E_+^2 - m_+^2) - (E_-^2 - m_-^2)}{2eBs} \quad (4)$$

The results quoted are from Eq. 2.

2. Results and discussion

Fig. 1 displays a typical mass shift as a function of the field for the proton. There is good linear behavior going through the origin, an indication that contamination from the higher terms has been effectively eliminated. We only use the two smallest field values to do the linear fit. Fig. 2 shows the results for the proton and the neutron. The line is a simple chiral fit using the ansatz $\mu = a_0 + a_1 m_{\pi}$. A more advanced chiral extrapolation could be carried out, using, for example, the prescription in Ref. [9,10]. Fig. 3 to Fig. 7 show the results for other particles. The agreement with experiment for the Ω^- is encouraging since no extrapolation is needed for this point. The experimental value for Δ^+ (slightly shifted for better view) is taken from Ref. [11]. Finally, Fig. 8 shows the proton and Δ^+ together. The opposite curvatures are a signature of quenched chiral physics. A similar behavior has been observed by [12,13] using the form factor method.

This work is supported in part by U.S. Department of Energy under grant DE-FG02-95ER40907, and by NSF grant 0070836. The computing resources at NERSC and JLab have been used.




Figure 1. Proton mass shifts as a function of magnetic field B at the six quark masses. The line is a fit using only the two smallest B values.




Figure 2. Magnetic moments (in nuclear magnetons) for the proton and neutron as a function of m_{π}^2 . The line is a simple chiral fit. The experimental values, taken from the PDG [8], are indicated by the empty symbols.









Figure 3. Magnetic moments for the octet Σ .

Figure 4. Magnetic moments for the octet Ξ .Figure 5. Magnetic moments for the Δ .Figure 6. Magnetic moments for the decuplet Σ^* .Figure 7. Magnetic moments for the decuplet Ξ^* .Figure 8. Comparison of magnetic moments of the proton and Δ^+ .

REFERENCES

1. Martinelli *et. al.*, Phys. Lett. **B116**, 434 (1982).
2. C. Bernard, T. Draper, K. Olynyk, Phys. Rev. Lett. **1076**, 269 (1982); Nucl. Phys. **B220**, 508 (1983).
3. J. Smit and J.C. Vink, Nucl. Phys. **B286**, 485 (1987).
4. H.R. Rubinstein, S. Solomon and T. Wittlich, Nucl. Phys. **B457**, 577 (1995).
5. J. Christensen, W. Wilcox, F.X. Lee, L. Zhou, hep-lat/0408024.
6. L. Zhou, F.X. Lee, W. Wilcox, J. Christensen, hep-lat/0209128.
7. L. Zhou, PhD dissertation (2004) (unpublished).
8. Particle Data Group, Phys. Lett. **B592**, 1 (2004).
9. R.D. Young, D.B. Leinweber, A.W. Thomas, hep-lat/0406001.
10. D.B. Leinweber, Phys. Rev. **D69**, 014005 (2004).
11. M. Kotulla *et. al.*, Phys. Rev. Lett. **89**, 272001 (2002).
12. J. Zanotti, D.B. Leinweber, A.W. Williams, J.B. Zhang, Nucl. Phys. B (Proc. Suppl.) 129-130, 287 (2004), hep-lat/0309186.
13. R.D. Young, D.B. Leinweber, A.W. Thomas, Nucl. Phys. B (Proc. Suppl.) 129-130, 290 (2004), hep-lat/0309187, hep-lat/0311038.